

THE EFFECT OF NATURAL ZEOLITE ON FATTENING LAMBS PRODUCTION RESULTS

Jovan Stojkovic^{1*}, Zoran Ilic¹, Milinko Milenkovic¹, Boban Jasovic¹, Goce Cilev²

¹**Faculty of Agriculture, University of Prishtina, Jelena Anzujska bb, Zubin potok, 38228, Kosovska Mitrovica, Republic of Serbia**

²**Veterinary Faculty, University St. Kliment Ohridski, Prilepska bb, 7000 Bitola, Republic of Macedonia**

*e-mail: jovan.s@jotel.co.rs

Abstract

The paper presents results of the investigation concerning the effect of a minazel-based preparation on fattening lambs production results.

The experiment was performed on two groups of lambs (control – K and experimental – O), 20 animals per group, and for a duration of 90 days. The diet consisted of sheep milk, a compound mix for fattening lambs, and meadow hay. Lambs of the experimental group, contrary to those of the control group, got a minazel-based preparation. The preparation Min-a-Zel S (in the form of 25% suspension) was administered, to lambs from birth until the 14th days of life, directly into the mouth, once daily (before the morning meal), 10 mL each. Min-a-Zel Plus was administered starting from the 15th day by adding it to the fodder mix (0.25%). Feeding was ad libidum.

The average starting lamb body mass was 4.10 kg. The average body mass of lambs at the end of the experiment was 24.50 vs 28.10 kg, K vs O respectively ($P < 0.01$). The daily grow of the lambs, during the course of the experiment, was 226:266 g, and was 40 g or 11.76% higher in the experimental group ($P < 0.01$). The experimental group had better utilization of dry matter, proteins and energy, with a lower incidence of diarrhoea.

Key words: Lambs, growth, feed, utilization, natural zeolite.

1. Introduction

The contemporary cattle production demands constant invention of the new possibilities that would improve health condition and production results of the domestic animals. Having in mind that aim, during the last decade a large number of reports dealing with possible use of natural zeolite in the cattle production could be met in public. Hence, it is believed that because

of its good absorption efficiency, the absorption of various harmful material can remove some bad or favorite influence of some good factors in feed animal digestive tract, as in the environmental conditions, under which the animals are kept Adamović *et al.* [2].

Zeolites are cationic ionizer containing clinoptilolite. They are mostly used for absorption of mycotoxins, i.e. for preventing of mycotoxicosis development in the animals. For this purpose 0.2-5% is added in the fodder mix. Previous investigation of Stojković *et al.* [7] showed the positive effects of using the preparation based on natural zeolite in the fattening lamb diet. This investigation is an addition to the previous one.

2. Materials and Methods

The experiment was performed on the two lamb groups (control - K and experimental - O), with 20 animals per group, at the farm Mramor near Nis. The experiment lasted 90 days. The feeding was ad libidum. The preparation Min-a-Zel S (in the form of 25% suspension) was administered to lambs from birth until the 14th day of life, directly into the mouth. Min-a-Zel Plus was administered starting from the 15th day by adding it to the fodder mix (0,5%). Min-a-Zel S is 25% suspension of refined zeolite compound mix, which contains 90% of clinoptilolite. Min-a-Zel Plus represents a new generation of mineral mycotoxin absorbents, and it is made by modification of the surface of zeolite material named clinoptilolite with a long-chained quaternary amine (Tomašević-Čanović Magalena *et al.* [8]). Both preparations were made at The Institute for Technology of Nuclear and other Mineral Raw Materials in Belgrade.

At the start of this experiment the usual chemical analyses of the feed components (Table 1) were made (milk, concentrate fodder mixture, and hay).

Table 1. Chemical composition of used feedstuffs (calculated to 100% of DM)

Parameter	Concentrate mixture	Meadow hay	Sheep milk
Dry matter	88.00	89.20	18.60
Proteins	17.16	11.88	32.26
Ash	6.36	7.62	3.49
Fat	3.08	2.24	39.52
Cellulose	11.02	30.72	0.00
NEF	62.39	47.56	24.73
Ca (g)	0.82	0.76	1.13
P (g)	0.64	0.56	0.81
ME, MJ/kg	8.07	5.11	13.45

The body mass of lambs was measured at the beginning and at the end of the experiment. The amount of consumed food was measured every day. The statistical interpretation of the data was done by Statistica computer program, version 6, StatSoft. Inc [6].

3. Results and Discussion

The results of lamb body mass gain indicate that lambs that received Min-a-Zel S and Min-a-Zel Plus gained more weight. The average body mass of lambs at the end of the experiment (90th day of life), was 24.50 vs. 28.10 kg (K vs. O, respectively), and the observed differences were statistically significant at the level of $P < 0.01$. The average lamb daily growth (Table 2) during the experiment was 226 vs. 266 g, and the established difference of 40 g or 11.76% in favor of the experimental group, was highly significant ($P < 0.01$). The difference in daily growth, in favor of the experimental group, was greater in the second part of the experiment, when lambs took most of the nutritive material from the concentrate fodder mixture in which Min-a-Zel Plus was added. The characteristics of the new generation preparation, enable it to bind large number of adverse materials (mycotoxins, radionuclides, heavy metals, ammonia, etc.) which can come into the organism through food, preventing their bad effects thus.

Table 2. The results of the comparative examination of natural zeolite in fattening lambs diet

Parameter	Group		Index
	K	O	
Number of lambs	20	20	-
Duration of the experiment, days	90	90	-
Input lamb body mass, kg	4.10	4.10	-
Output lamb body mass, kg	24.50	28.10**	114.89
Total growth of lambs, kg	20.40	24.00**	117.64
Daily growth of lambs, g	226	266**	117.69
Food consumption, kg	2.40	2.60*	108.33
Food conversion, kg	9.20	8.60*	93.47

* $P < 0.05$; ** $P < 0.01$

Previous reports also point out to positive effects of minazel on the production results of fattening lambs.

In the previous experiment performed by Masić *et al.* [4] the body mass of the lambs in the control group at the end of the fattening was 24.48 kg, while in the experimental group, with added preparation based on zeolite, was 26.94 kg. Abu-Zanat [1] reached daily growth from 167 to 197 g by added clinoptilolite in lamb meals. Using the same zeolite-based preparation (Min-a-Zel Plus) in fattening lambs in dose of 0.2% and 0.5% of concentrated mixture Adamović *et al.* [3] observed the daily growth of 209 g and 179 g respectively, while the increased preparation dose did not increase daily growth additionally. However, in this research an increased dose of Min-a-Zel Plus influenced zearalenon residue elimination from liver, kidneys, and muscles, what was the real importance of these results. Adding in mixture for older lambs 0.2 or 0.5% of the zeolite-based preparation (Min-a-Zel), Stojković *et al.* [7] reached slight increase of lamb body mass (2%), and daily growth (3%), what was significantly less than the daily growth increase found in this research Abu-Zanat [1] also affirmed that the lambs eating the mixture with clinoptilolite added showed better results, while Pupavac *et al.* [5] did not observe any positive result using lamb diet with added zeolite, which probably was the result of using different zeolite type.

The lambs food consumption, according to the sequence of treatments K vs O was 2.40 vs 2.60 kg respectively, and the observed difference was statistically significant ($P < 0.05$). The results of the food conversion indicate that the lambs fed by additional Min-a-Zel S and Min-a-Zel Plus spent less food per one kg growth. During the whole experiment (0-90 days), the food conversion of the studied groups K vs. O was 3.20 vs. 3.00 kg respectively, and the observed difference was highly significant ($P < 0.05$).

A greater difference in growth observed in this study probably is a result of these two zeolite-based preparations combined influence in the different stages of growth. Application of Min-a-Zel S suspension (used in the first 14 days of life) contributed to the better results of experimental group during the first weeks of lambs growth, when the use of the concentrate mixture was insignificant. Min-a-Zel Plus preparation had the positive influence after 30th day, when the lambs' became used to eat a larger amount of the mixture in which this preparation was added.

In the control group lambs between a 17th and 23rd day of life, seven lambs had diarrhoea, which repeated between 45th and 50th day of life. In the experimental group lambs any serious diarrhoea case was not noticed, which probably was a contribution of the tested preparations.

4. Conclusions

- The use of the preparation based of natural zeolite (Min-a-Zel S and Min-a-Zel Plus) had a positive influence on the production of the fattened lambs.
- The average body mass of the control group lambs was 24.50, while the average body mass of the experimental group lambs was 28.10 kg, which was by 3.60 kg or 11.48% more.
- The average daily growth was greater in the experimental group for 40 g or 11.76%. In the same group the food consumption per kilogram of growth was by 60 g or 6.53%.
- The diarrhea incidence rate was also lower in the experimental group.

5. References

- [1] Abu-Zanat M.M.W. (1998). *Using natural zeolites in the fattening of Awassi lambs*. Dirasat-Agriculturale-Scienes, 24, (2), 268-273.
- [2] Adamović M., Nešić S., Stpoičević S., Tomašević-Čanović Magdalena. (2001). *Uticao organski modifikovanog adsorbenta Min-a-Zel Plus na kvalitet silaže biljke kukuruza*. IX Jugoslavenski simpozijum o krmnom bilju. Beograd.
- [3] Adamović M., Tomašević-Čanović Magdalena, Milošević S. (2002). *Doprinos mineralnih adsorbenata mikotoksina povećanju proizvodnje i kvalitetu hrane*. Eko-Konferencija, Novi Sad.
- [4] Mašić Z., Kljajić R., Škrinjar M. (1999). *Mikotoksini u stočnoj hrani kao faktor poremećaja zdravlja životinja*. 12 Savetovanje veterinarara, 65-73, Vrnjačka banja.
- [5] Pupavac S., Sinovec Z., Nešić S., Hudina V., Stevanović J. (2001). *Uticao korišćenja različitih adsorbenata mikotoksina u ishrani prasadi*. Arhiv za poljoprivredne nauke, 61, 193-103. Beograd.
- [6] StatSoft. Inc. *Statistica, verzion 6*.
- [7] Stojković J., Sinovec Z., Adamović M., Tomašević-Čanović Magdalena. (2004). *Efikasnost primene preparata na bazi zeolita u obrocima jagnjadi u tovu*. Vet. Glasnik, Beograd, Vol. 58, br. 3-4, 327-334.
- [8] Tomašević-Čanović Magalena, Dumić M., Olivera Vukičević, Aleksandra Daković, Rajić I. (2000). *PATENT P-838/2000*.